

# Annual WWW Technical Progress Report

## On the Global Data Processing and Forecasting System 2004

### FRANCE

#### 1. SUMMARY OF HIGHLIGHTS

- . enhanced configuration of the supercomputer Fujitsu VPP5000 with 124 processors
- . use of ATOVS AMSU-B radiances from NOAA satellites and Quikscat data in ARPEGE analysis
- . short range ensemble forecast using ARPEGE global model

#### 2. HARDWARE USED

- Information commutators on GTS are the TRANSMET computers (2 Sun Enterprise 3000, operating with OS Unix and RDBMS Oracle).
- the management of the forecasting system (control of the data in input of NWP models, postprocessing, production of charts with the NWP output) is made on a HP N4000 computer running Oracle RDBMS, US-Navy originating NEONS meteorological data management system, and PV-WAVE graphical software; one HP K580 is used as file server, one HP C180 workstation is devoted to the system monitoring, which is based on DCE. The whole system (production machine + file server + monitoring workstation), called DIAPASON, is doubled for backup.
- NWP operational models are running on a FUJITSU VPP5000 (31 processors, 21 with 8 Gbytes memory each, the others with 4Gbytes memory each)
- Dissemination of forecast and observation products (from GTS included), in particular to the french weather stations, is made through satellite communication (RETIM2000 system).

#### 3. USE OF DATA AND PRODUCTS FROM GTS

Average number of messages, by day:

AIREP	ACARS	AMDAR	BATHY	BUOY	PILOT	SATEM	SATOB	SHIP	SYNOP	TEMP	TEMPSH IP
3000	130000	24000	40	14000	1700	20000	75000	12000	60000	1300	17
HIRS	AMSU-A	AMSU-B	Sea wind	ERS URA	ERS UWA	ERS UWI	PROFIL ER-US	PROFIL ER-EU	GEO RAD	GEO WIND	SSMI
43000	3200	100000	25000	180	250	250	750	2000	4500	5000	46000

- GRID from EXETER : 2876
- GRID from WASHINGTON : 525
- GRIB aero 1.25 from EXETER : 18816
- GRIB 2.5 from EXETER : 9552
- GRIB ECMWF : 1776
- Fac-simile products:
  - aeronautical charts from Exeter 476 and ECMWF 110 (T4 code)

#### 4. DATA INPUT SYSTEM

Automated.

## **5. QUALITY CONTROL SYSTEM**

GTS data are controlled at several levels:

- transmission
  - syntactic coherence
  - rudimentary control of likelihood: e.g. a sea level pressure value must be above 880 hPa and below 1080 hPa
  - data control by comparison to adjacent (in time and/or space) data, or to different types of data at the same location: e.g. Td T is checked; In the same manner, a sudden slope breaking of in a temperature profile from a radiosonde far from the tropopause leads to the invalidation of the data.
- Data to be reemitted on GTS are not modified.

## **6. MONITORING OF THE OBSERVATION SYSTEM**

All the observations that are used by the NWP system (SYNOP, SHIP, BUOY, SATOB, ATOVS, TEMP, PILOT, AIREP, AMDAR, ACARS) are controlled by comparison to the analyses and first guesses of the ARPEGE assimilation cycle: statistics are produced every month and summarized in a monthly bulletin.

## **7. FORECAST SYSTEM**

The operational forecast system at Météo-France is based on two different numerical applications of the same code ARPEGE-IFS and an additional code to build the limited area model ALADIN.

The ARPEGE-IFS library has been developed jointly by Météo-France and ECMWF (ARPEGE being the usual name in Toulouse and IFS the one used in Reading):

ECMWF model for medium range forecasts (4-7 days)

a variable mesh version run in Toulouse for short range predictions (1-4 days)

The ALADIN library has been developed jointly by Météo-France and the national meteorological or hydrometeorological services of the following countries: Austria, Belgium, Bulgaria, Croatia, Czech Republic, Hungary, Moldova, Morocco, Poland, Portugal, Romania, Slovakia, Slovenia, Tunisia.

### **7.1. Schedule of the Forecast System**

The operational forecast system at Météo-France is based on ARPEGE/ALADIN, using the following rules :

. an assimilation (long cut-off) analysis is performed before each short cut-off analysis.

. the product's availability is :

initialised analysis (P0)	cut-off+30'
ARPEGE forecast	20' every 24H range
ALADIN-France	ARPEGE+10'

Stretched configuration

HH	0000 UTC	0600 UTC	1200 UTC	1800 UTC
long cut-off	0810 UTC	1250 UTC	2010 UTC	0050 UTC
short cut-off	1H50	3H	1H50	3H
ARPEGE range	102H	72H	72H	60H
End of ARPEGE	0340 UTC	1010 UTC	1520 UTC	2200 UTC
ALADIN range	54H	48H	42H	36H
end of ALADIN	0310 UTC	1020UTC	2300 UTC	2210UTC

### **7.2. Medium range (4-10 days) forecast system**

As mentioned above, it is the operational T511 IFS model of ECMWF and T255 Ensemble Prediction System for 4-5day and 6-7day forecast bulletins.

### **7.3. Short range forecast system**

#### **The ARPEGE system(0-102 hours)**

ARPEGE-IFS is a common Météo-France / ECMWF development. ARPEGE is the French name (Action de Recherche Petite Echelle Grande Echelle) while IFS is the

name used at Reading (Integrated Forecast System). It is a tunable system based on a global spectral model which can be used for several applications: data assimilation, short-range prediction, medium-range prediction, climate research, predictability studies.

ARPEGE-IFS uses Schmidt's transformation leading to variable mesh configurations, having a pole of maximum resolution and a resolution varying continuously from that pole to the antipode (Courtier and Geleyn 1988). T being the nominal truncation and C the "stretching factor", the local resolution of the model is T x C over the pole, and T / C at the antipode.

The present version is T358 C2.4 having its grid pole in France (46.5N,2.6E), leading to a horizontal resolution of the linear grid of 23 km over France and 133km over New Zealand.

The number of vertical levels is 41, with an increased density in the low atmosphere. The first level is at 1 hPa, and the lowest one at 17m above the ground.

### **Assimilation, objective analysis and initialization**

The assimilation runs with a 6 hour cycle. The objective analysis is performed with a multi-incremental 4D variational scheme : i.e. the departure obs-guess is computed at full resolution (T358C2.4) whereas the analyzed structures are produced at a lower resolution, in 2 loops T107C1, T149C1. It is therefore assumed that the small scales (not corrected by the analysis) are forced by the (analyzed) large scales in the subsequent forecast.

<b>assimilated data:</b>	SYNOP, SHIP, BUOY, BATHY, TEMP, TEMPSHIP and PILOT (part A, B, C and D), profilers, AIREP, AMDAR, ACARS, SATOB, ATOVS (AMSU-A, B and HIRS), Quikscat ambiguous winds, with observation time in [H-3h,H+3h]
<b>assimilation cycle:</b>	6 hour cycle.
<b>analysis method:</b>	Multivariate four dimensional variational analysis
<b>analysed variables:</b>	Wind, temperature and specific humidity on model levels, plus surface pressure.
<b>first guess:</b>	A 6-hour forecast of ARPEGE in normal operations
<b>cover:</b>	Global cover.
<b>horizontal resolution:</b>	T149 linear grid increments on a T358C2.4 background
<b>vertical resolution:</b>	The analysis is done on the model levels (see below): 41 levels (hybrid vertical co-ordinate) from 18m up to 1 hPa.
<b>initialization:</b>	weak DFI constraint in the variational cost function and incremental digital filter initialization (ie filtering analysis increments fields) using a Dolph-Chebichev filter with a stop-band edge period of 5h .
<b>surface:</b>	-analysis of superficial and mean soil temperature (resp moisture) from forecast errors on 2m temperature (resp. relative humidity)relaxation towards climatology for snow and mean soil temperature and moisture initialisation of Sea Surface Temperature and ice concentration using NOAA data

### **Model**

<b>basis equations:</b>	Primitive equations system
<b>independent variables:</b>	horizontal wind vector, temperature, specific humidity and surface pressure.
<b>dependent variables:</b>	Vertical velocity and density
<b>numerical technique:</b>	Spectral two-time-level semi-lagrangian model and temporal discretization using semi-implicit scheme
<b>integration domain:</b>	global

<b>orography, gravity wave drag:</b>	The orography of this model is computed on the T358 C2.4 Gaussian grid (358x720 points) from GLOB95 30"+US NAVY 10' + NOAA 5' data using a variational technique that strongly reduces the noise associated to Gibbs waves (see Bouteloup, 1995). The gravity wave drag takes in account some anisotropy, blocking and mid-tropospheric effects.
<b>horizontal diffusion:</b>	Implicit in spectral space and incorporating an orography dependent correction for temperature
<b>vertical diffusion:</b>	Scheme linked to PBL (see next point)
<b>planetary boundary layer:</b>	ECMWF method (Louis et al. 1981) with several enhancements in the stable case
<b>resolution, time step:</b>	This version of the ARPEGE model has a triangular linear grid type truncation T358 with a stretching factor C2.4. It has 41 vertical levels from 18m up to 1hPa, using the hybrid (s,p) co-ordinate from Simmons and Burridge (1981). The time step is 16 minutes.
<b>Earth surface:</b>	An improved version of the ISBA (Interaction Soil Biosphere Atmosphere) scheme is used, including an explicit parameterization of soil freezing. Six prognostic variables are handled by ISBA: surface temperature, mean soil temperature, interception water content (water on the leaves), superficial soil water content (first centimeter), total liquid soil water content, total frozen soil water content. A very simple parameterization of snow cover is added with albedo ageing. Soil characteristics (texture, depth) are point-dependent. Vegetation characteristics are point- and month-dependent.
<b>radiation:</b>	Highly simplified scheme (inspired from Ritter and Geleyn 1992) called at every time-step in every grid-point. Solar terms, cooling-to-space and exchange-with-surface terms for the thermal range are computed exactly but with one single spectral interval, exchange-between-layers thermal terms are further approximated by a "no-overestimation" simplification of the saturation process.
<b>convection:</b>	Mass-flux scheme (Bougeault 1985) enhanced with <ul style="list-style-type: none"> <li>i. the Gregory-Kershaw treatment of momentum transport by cumulus,</li> <li>ii. a treatment of the moist adiabatic computation consistent with "i",</li> <li>iii. a downdraft parameterisation,</li> <li>iv. vertically variable entrainment and detrainment rates,</li> <li>v. a parameterisation of the selective effect of entrainment leading to a warmer upper part of the single cloud ascent.</li> </ul>
<b>humidity:</b>	Specific humidity is the variable: no storage of condensate; evaporation of falling rain; treatment of the ice-phase.

### **PEARP ensemble forecasting system (0-60 hours)**

This is an ensemble of 10 perturbed forecasts in the vicinity of the 18UTC ARPEGE analysis, run using the same ARPEGE model as above. The perturbation method is based on T95L41 24-hours singular vectors targeted on Western Europe.

### **ALADIN (0-54hours)**

ALADIN is a limited area version of ARPEGE-IFS.

- ALADIN is spectral
- As a spectral it works on a bi-periodic domain and uses bi-Fourier horizontal transforms

- Same physics as ARPEGE
- Initial and lateral boundary conditions from ARPEGE

ALADIN is run in pure dynamical adaptation mode, i.e. without own data assimilation. The operational version is semi-lagrangian (timestep 7min), with elliptic linear grid type truncation E149x149 on Lambert projection domain (54°95N/33°66N,-11°18W/19°64E), leading to an equivalent finite difference resolution of roughly 9km.

The vertical resolution is 41 levels, the same as operational ARPEGE ones. The digital filter initialization uses a Dolph-Chebyshev filter with a stop-band edge period of 3h and a backward-forward scheme.

### **NWP Products**

The above described numerical models feed a analysis and forecast database, having following characteristics:

- different horizontal domains for different horizontal resolution (from the global domain with a 2.5° and 1.5° mesh to the "France" domain with a 0.1° mesh)
- vertical levels are the standard pressure levels
- independence, from the creating model, of the format of the database products.

The meteorological fields stored in this database are:

- at all levels: geopotential, temperature, humidity, wind (including vertical velocity)
- at screen level: pressure, temperature, humidity, heat and radiation fluxes, snow and water content
- at sea surface level: reduced pressure
- some data at particular levels: 500 hPa absolute vorticity, high medium and low cloudness, iso 0° and iso -10°, tropopause etc...

ARPEGE produces boundary conditions for the ALADIN applications run in Austria, Bulgaria, Croatia, Czech Republic, Hungary, Morocco, Poland, Portugal, Romania, Slovakia, Slovenia, Tunisia, while ALADIN-France provides boundary conditions for ALADIN-Belgique.

### **Operational use of NWP products**

On screen (especially SYNERGIE workstation and Meteotel-PC software) or on paper, hundreds of charts...

## **7.4. Specialized forecasts**

### **7.4.1 Local weather elements**

Millions of local forecasts of weather parameters are produced daily through statistical adaptation of NWP output. Main methods are multiple linear regression (MLR) and discriminant analysis (DA). MOS (model output statistics) is preferred to PP (perfect prognosis). Kalman filter (KF) is applied when relevant. Ensemble distributions are calibrated before computing probabilities.

- 2m temperature: MLR+KF  
ARPEGE model, France: 1182 stations. +3h to +96h (resp 42h, 72h and 30h) by 3h from 00 UTC (resp.06 UTC, 12 UTC and 18UTC) + daily extremes.  
ECMWF model, France: 1182 stations. +12h to +180h by 3h from 12UTC + daily extremes.  
ECMWF model, world: 6010 stations. +6h to +180h by 3h + daily extremes.
- 2m temperature: MLR applied to individual runs, KF applied to ensemble mean.  
ECMWF EPS, France: 206 stations. Daily extremes day+1 to day+8.
- 2m temperature, probabilities: MLR applied to individual runs, ensemble calibration.  
ECMWF EPS, France: 1153 stations. Daily extremes day+1 to day+8.
- 2m humidity: MLR+KF  
ARPEGE model, France: 990 stations. +3h to +96h (resp 42h, 72h and 30h) by 3h from 00UTC (resp.06 UTC,12 UTC and 18UTC).

- Dew point temperature: from temperature and humidity products  
ARPEGE model, France: 982 stations. +3h to +96h (resp 42h, 72h and 30h) by 3h from 00UTC (resp.06 UTC,12 UTC and 18UTC).
- Total cloud cover: MLR+KF  
ARPEGE model, France: 139 stations. +3h to +96h (+72h) by 3h from 00UTC (12UTC).  
ECMWF model, France: 139 stations. +12h to +240h by 3h from 12UTC.
- Total cloud cover, probabilities (categories 0, 0-2, 1-2, 3-4, 3-5, 5-6, 5-7, 6-8, 7-8, 8): DA  
ARPEGE model, France: 150 stations. +3h to +96h (resp 42, 72 and 30h) by 3h from 00UTC (resp.06 UTC, 12 UTC and 18UTC).
- 10m wind speed: MLR  
ARPEGE model, France: 645 stations. +3h to +96h (+72h) by 3h from 00UTC (12UTC).
- 10m wind speed, probabilities: MLR to individual runs, ensemble calibration.  
ECMWF EPS, France: 213 stations. +24h to +240h by 24h from 12 UTC.
- Visibility, probabilities (thresholds 800, 1000, 1500, 3000 and 5000m):  
DA  
ARPEGE model, France: 16 stations. +18H from 12UTC.
- Wind gusts, probabilities (thresholds 28 to 78kt by 5kt): DA  
ARPEGE model, France: 373 stations. +3h to +96h (resp 42h, 72h and 30h) by 3h from 00UTC (resp 06 UTC,12 UTC and 18 UTC).
- 24h precipitations, probabilities: ensemble calibration.  
ECMWF EPS, France: 776 stations. day+1 to day+8 from 12 UTC.

#### **7.4.2 Marine forecasts**

##### **Wave hindcast and forecasting system**

Four models run operationally in France for determining the sea conditions:

**A global wave model** , computing the waves over all the oceans up to 102 hour forecast, from the wind outputs of large scale fields derived the global atmospheric models ARPEGE

Type:	coupled discrete deep water
Integration domain:	Global
Grid:	regular grid; resolution: 1°
Frequency resolution:	12 frequency components, logarithmically spaced from 0.04 Hz to 0.3 Hz
Direction resolution:	18 equally-spaced direction components
Integration scheme:	time step = 900s
Boundary forcing:	winds at 10m level from ARPEGE, updated every 6 hours
Surface classification:	sea ice deduced from ARPEGE SST
Assimilation:	4 analysis/day using significant wave heights from ERS2 altimeter

**Another global wave model** , computing the waves over all the oceans up to 72 hour forecast, from the wind outputs of large scale fields derived the global atmospheric models ARPEGE/TROPIQUE

Type:	coupled discrete deep water
Integration domain:	Global
Grid:	regular grid; resolution: 1°
Frequency resolution:	12 frequency components, logarithmically spaced from 0.04 Hz to 0.3 Hz
Direction resolution:	18 equally-spaced direction components
Integration scheme:	time step = 900s
Boundary forcing:	winds at 10m level from ARPEGE/TROPIQUE, updated every 6 hours
Surface classification:	sea ice deduced from ARPEGE SST
Assimilation:	4 analysis/day using significant wave heights from ERS2 altimeter

**A regional model**, forecasting the waves up 54 hours with 3 hour step, over the European Seas (Atlantic, Mediterrean , Baltic, North Sea, Black sea, ...) , from the wind outputs of small scale fields derived from ARPEGE.

Type:	Coupled discrete shallow water
Domain:	European Seas : 67N-30N-12W-42E
Grid:	regular grid; resolution: 0°25
Frequency resolution:	12 frequency components, logarithmically spaced from 0.04 Hz to 0.3 Hz
Direction resolution:	18 equally-spaced direction components
Timestep:	300s
Boundary forcing:	winds at 10m level from ARPEGE, updated every 3 hours.

**A coastal model**, forecasting the waves up 54 hours with 3 hour step, over the French contiental shelf, from the wind outputs of small scale fields derived from ALADIN.

Type:	Coupled discrete shallow water
Domain:	French Seas (Metropolitan France only): 57N-35N-11W-17E
Grid:	regular grid; resolution: 0°1
Frequency resolution:	12 frequency components, logarithmically spaced from 0.04 Hz to 0.3 Hz
Direction resolution:	18 equally-spaced direction components
Timestep:	150s
Boundary forcing:	winds at 10m level from ALADIN, updated every 3 hours.

These models are available on 00UTC, 06UTC, 12UTC and 18UTC runs, except for VAG/TROPIQUE, available only on 00UTC and 12UTC.

#### **Operational simulations of the oceanic circulation in tropical Atlantic**

The oceanic primitive equation model OPA7, developed by CNRS/LODYC, has been run operationally every month, using all the surface fluxes produced by the operational ARPEGE model. Its main characteristics are 17 horizontal levels in z coordinate with a realistic bathymetry, and a 1/3 degree horizontal resolution. Systematics comparisons have been performed with bathythermic observations sent through the GTS, and against sea surface temperatures from ERS data ( ATSR ).

#### **Storm surge model**

A depth-averaged, numerical storm-surge model has been developed and configured to provide storm-surges forecasts along coastlines of France. Two versions of this model, one for overseas territories to forecast tropical cyclones storm surges and one for metropolitan French coastline.

**Overseas domain:** Atmospheric fields are inferred from an analytical-empirical cyclone model which require only cyclone position, intensity and size. The model has been operated since 1994 in the French Antilles, 1995 in New Caledonia, 1997 in the French Polynesia and 1998 in La Reunion. The model can be used in two different ways. In real-time mode as a tropical cyclone is approaching an island or in climatological mode: a cyclone climatology is used to prepare a data base of pre-computed surges. Due to the low accuracy of tropical cyclone trajectory forecasts, the second mode seems to be, at present time, the best way to use the model.

The grid mesh is fixed for each domain and varies from 150 m to 1850 m.

**Metropolitan domain:** Atmospheric fields are taken from atmospheric numerical models: IFS (ECMWF), ARPEGE and ALADIN (Météo-France). The system has been operated since October 1999 for the Channel and Bay of Biscay, March 2002 for the Mediterranean Sea and November 2002 for the North Sea.

48 hours forecast are produced on a 5' grid mesh.

### **Drift model (oil spills, containers, Search & Rescue)**

Météo-France is in charge of spill drift predictions within the spill response plan POLMAR-MER in case of a threat for the French coastline. At an international level, Météo-France can intervene within the Marine Pollution Emergency Response Support System (MPERSS) for the high seas. Météo-France is Area Meteorological Co-ordinator for METAREA II and III west, and supporting service for METAREA I, III east, VII B and VIII C.

Météo-France developed a drift model named MOTHY (Modèle Océanique de Transport d'HYdrocarbures). MOTHY is an integrated system that includes hydrodynamic coastal ocean modelling (2D+1D) and atmospheric forcing from ARPEGE or IFS models. The hydrodynamic coastal ocean is linked to an oil spill model, where oil slick is considered as a distribution of independent droplets. These droplets move with shear current, turbulent diffusion and buoyancy. The system has been operated since 1994 and can be used for oil spills or drifting objects. New developments, exercises and training are jointly conducted with CEDRE (Centre de documentation de recherche et d'expérimentations sur les pollutions accidentelles des eaux). MOTHY correctly predicted the drift of the oil during Erika (December 1999) and Prestige (2002-2003) in the Bay of Biscay.

The domain is global with a better accuracy on specific areas, including French seas. Forecasts are produced up to 5 days on fixed grid from 150 m to 9 km.

### **7.4.3 Pollutant transport and dispersion forecast**

At an international level, Meteo-France Toulouse has been designated as a regional specialized meteorological centre (RSMC) with activity specialization on the provision of atmospheric transport model products for environmental emergency response. This provision is related to nuclear accident, or radiological emergencies, and plumes of volcanic ashes for ICCA. In the framework of the french government emergency plan, Meteo-France is also involved for chemical or nuclear releases. The operational organization of Météo-France, for facing atmospheric pollution accidents, is based on a special crisis meteorological cell (CMC) that studies the evolution of weather/pollution conditions and provides forecasts on the pollutant plume.

Up till now, for the long-range dispersion forecast, Meteo-France Toulouse uses two tools to track the plume following an accidental release: an air mass trajectories software, describing the evolution of a neutrally buoyant particle in the wind field forecasted by the NWP, and an eulerian off-line dispersion model, MEDIA, solving an advection-diffusion equation for a passive scalar. A next generation of model will be implemented in 2005, MOCAGE-Accident. Indeed, Meteo-France has developed a global three-dimensional Chemical Transport Model, MOCAGE, dedicated to the numerical simulation of the interactions between dynamical, physical and chemical processes in the lower stratosphere and in the troposphere, for air quality forecasts. This model will be used in the specific way of a ponctual emission, with the set of sinks, as dry deposition, scavenging of soluble gases by convective and stratiform rain, and fallout. MOCAGE-Accident is currently evaluated with the intercomparison exercise ENSEMBLE (<http://ensemble.ei.jrc.it/>).

At local scale, the system PERLE focuses on the local description of the atmospheric pollutant cloud at regional and local scale, in the vicinity of the affected site of radionuclide or chemical release. It is based on a meso-scale non hydrostatic model for meteorological fields, Meso-NH (Lafore et al., 1998), coupled to a lagrangian particle model for the dispersion.

Meso-NH uses two nested models for emergency response, with a first domain covering 240km\*240km area (8-km resolution) and a second domain covering 60km\*60km area (2-km resolution), and two-way interactions between them. The initial and boundary conditions of the larger domain are defined by ALADIN. With advanced physical subgrid parametrizations (turbulence, convection ...), Meso-NH is more satisfactory than diagnostic models, based on the principle of conservation of mass, that are commonly used for dispersion modelling. Furthermore, passive tracer is simulated by the eulerian model, to provide a regional description of the pollutant cloud at regional scale. The

lagrangian particle formulation allows a description of the pollutant cloud in the vicinity of the release (30km\*30km area) during the first critical few hours, without gaussian assumptions commonly used at this scale.

#### **7.4.4 Tropical cyclones forecast model**

A specific version of ARPEGE, called ARPEGE-Tropiques, has been implemented for more detailed forecasts over tropical areas, and sent to the SYNERGIE software in French oversea regional centers.

The ARPEGE-Tropiques model is the same as the metropolitan one, but with a uniform truncature T358L41 for the forecast (time step 1350s) , and T107C1L41 for the 4DVAR analysis. Sea surface pressure bogus data , produced by forecasters in La Réunion center are incorporated in the assimilation to get a more precise location of cyclones. These bogus data are transmitted on the GTS in BUFR.

The models are running once a day based on 00UTC, up to 72 h , with a 3.50 hour cut-off.

Uniform configuration

HH	0000 UTC	0600 UTC	1200 UTC	1800 UTC
long cut-off	0955 UTC	2145 UTC	2210 UTC	0200 UTC
short cut-off	3H50			
ARPEGE range	72H			

#### **7.4.5 Snow and avalanches**

For several years, applications related to snow and avalanches in Grenoble have used the ensemble of models "SAFRAN / CROCUS /MEPRA". SAFRAN is an analysis system working at the scale of one mountain system (massif). The system has also been exported to various foreign countries.

Since the end of 2001, this new analysis system has been run operationally. It allows the use of surface observations with a 1hour frequency. It is used in forecast mode over the Alps and the Pyrenees, with precipitation fields from ALADIN/France as input.

#### **7.4.6 Hydrology**

The analysis code SAFRAN is also used for hydrological applications. In 2002 a version was installed in the operational departments of Météo-France, in order to produce precipitation estimates over France. In this hydrological mode SAFRAN is coupled to the surface scheme ISBA and to a hydrological model called MODCOU. The result of this coupling is known as SIM: SAFRAN – ISBA – MODCOU. The ARPEGE precipitations are used as input to the whole system.

### **7.6. Long range forecasts (3 months)**

A specific version of ARPEGE model , called ARPEGE-Climat is used 9 times a month to run 120 to 129 days forecasts, starting from ECMWF assimilation. The SST forecast is based on an auto regresssive statistical scheme on grid points, which is run once on the first day of the series. The seasonal forecasting system is using mainly the same ARPEGE software as the short range forecast model, except the following points:

- resolution, time step:** This version of the ARPEGE model has a triangular truncature T63 without stretching. The collocation grid has 128x64 points with a reduction near the poles; it has 31 vertical levels like IFS model during ERA-15 (ECMWF reanalysis). The time step is 1800 seconds.
- radiation:** Fouquart Morcrette scheme (1995)
- clouds, vertical diffusion, stratified precipitations:** Ricard Royer statistical scheme (1993).

## **8. VERIFICATION OF FORECASTS**

Scores of the operational ARPEGE model:

### **Against analyses**

	24 hours			72 hours		
	NH	SH	TR	NH	SH	TR
Z500 RMSE	10.9	15.7		30.3	40.9	
W250 RMSEV	4.9	5.1	4.8	10.4	11.3	8.1
W850 RMSEV			2.6			4.1

NH : Northern Hemisphere

SH : Southern Hemisphere

TR : Tropics

### **Against observations**

24 hours

	NA	EU	AS	AU/NZ	TR	NH	SH
Z500 RMSE	13.1	13.1	13.7	14.5	13.6	14.1	17.3
W250 RMSEV	6.9	6.1	6.5	6.3	6.0	6.4	6.6
W850 RMSEV	4.2	4.4	4.5	4.4	4.4	4.5	5.2

72 hours

	NA	EU	AS	AU/NZ	TR	NH	SH
Z500 RMSE	30.1	30.9	27.2	26.9	15.7	32.3	32.9
W250 RMSEV	12.1	11.2	10.4	10.1	8.1	11.3	10.9
W850 RMSEV	6.0	6.0	6.3	6.0	5.2	6.2	6.6

NA : North America EU : Europe AS : Asia AU/NZ : Australia / New Zealand

NH : Northern Hemisphere SH : Southern Hemisphere

TR : Tropics

### **Recall:**

Météo-France draws up a quarterly bulletin of "verification of the numerical products used for meteorological forecasting" (in French) which can be obtained by writing to:

Météo-France  
DPrévi/COMPAS  
42, av. Coriolis  
F-31057 TOULOUSE Cedex 1  
FRANCE

## **9. FUTURE PLANS**

For more details, see the Commission of Sciences / Progress Report on Numerical Weather Prediction.

- improvements to model physics (more sophisticated radiation, boundary layer mixing, cloud representation)
- data assimilation for the ALADIN regional model
- higher model top
- improved use of some existing observing systems (TEMP) and satellite data (AMSU, AIRS, Modis winds, Meteosat radiances)
- implementation of the coupled Arpege/Orca model at ECMWF and participation to the multi model operational experimentation with the ECMWF and Met Office models.

## REFERENCES

- Bougeault P., 1985 : "Parameterization of cumulus convection for GATE. A diagnostic and semi-prognostic study". *Mon. Wea. Rev.*, 113, 2108-2121.
- Bouteloup Y., 1995: "Improvement of the spectral representation of the earth topography with a variational method", *Mon. Wea. Rev.*, 123, 1560-1573
- Courtier, P. and J.F. Geleyn, 1988 : "A global numerical weather prediction model with variable resolution: application to the shallow-water equations". *Quart. J. Roy. Meteor. Soc.*, 114, 1321-1346.
- Giard, D., and E. Bazile, 1999 : Implementation of a new assimilation scheme for soil and surface variables in a global NWP model, submitted to *Mon. Wea. Rev.*.
- Louis J.F., 1979: "A parametric model of vertical eddy fluxes in the atmosphere", *Bound. Lay. Met.*, 17, 187-202
- Lafore, J.-P. et al., 1998 : The Meso-NH atmospheric simulation system. Part 1 : Adiabatic formulation and control simulations. *Ann. Geophysicae*, **16**, 209-228.
- Lynch, P., D. Giard and V. Ivanovici, 1997 : Improving the efficiency of a digital filtering scheme for diabatic initialization, *Mon. Wea. Rev.*, 125, 1976-1982
- Louis J.F., M. Tiedtke and J.F. Geleyn, 1981 : "A short history of the PBL parameterization at ECMWF". *ECMWF Workshop on PBL parameterization*, ECMWF, Reading, UK, 59-80.
- Morcrette, J.-J., 1991 : Radiation and clouds radiative properties in the European center for medium-range weather forecasts forecasting system. *J. Geophys. Res.*, 96, 9121-9132.
- Morcrette, J.-J., and Y. Fouquart, 1985: On systematic errors in parametrized calculations of longwave radiation transfer. *Quart. J. Roy. Meteor. Soc.*, 111, 691-708.
- Noilhan, J., and S. Planton, 1989 : A simple parameterization of land-surface processes for meteorological models, *Mon. Wea. Rev.*, 117, 536-549
- Piedelievre J.P., L. Musson-Genon and F. Bompay, 1990: MEDIA - An Eulerian model of atmospheric dispersion: first validation on the Chernobyl release, *Jour. of Appl. Met.*, vol. 29, N° 12, 1205-1220
- Ricard, J.-L., and J.-F. Royer, 1993: A statistical cloud scheme for use in an AGCM. *Ann. Geophys.* 11, 1095-1115
- Ritter B. and J.F. Geleyn, 1992 : "A comprehensive radiation scheme for numerical weather prediction models with potential applications in climate simulations". *Mon. Wea. Rev.*, 120, 303-325.
- Simmons A.J. and D.M. Burridge, 1981 : "An energy and angular momentum conserving vertical finite difference scheme on a hybrid vertical coordinate". *Mon. Wea. Rev.*, 109, 758-766.
- Yessad K. and P. Bénard, 1996 : "Introduction of a local mapping factor in the spectral part of the Météo\_france global variable mesh numerical model". *Quart. J. Roy. Meteor. Soc.*, 122, 1701-1719.